arXiv:physics/0012026v1 [physics.class-ph] 14 Dec 2000

In Memory of my Beloved Mama,
Vlasova Alexandra V.,
1922-2000

CHARGED BROWN PARTICLE: THE MORE RETARDATION IS - THE LOWER IS THE EFFECTIVE TEMPERATURE

Alexander A. Vlasov

High Energy and Quantum Theory
Department of Physics
Moscow State University
Moscow, 119899
Russia

The Brownian motion of a charged particle with finite size (described by Sommerfeld model) is considered. It is found out that due to radiation reaction: (1) the effective temperature of such particle is lower, and (2) the acceleration of the average velocity is smaller, then that for classical Brown particle without electric charge.

03.50.De

Sommerfeld particle [1] is the model of a charged particle of finite size - sphere with uniform surface charge Q and mechanical mass m. In nonrelativistic approximation such sphere obeys the equation (see [2] and works, cited there):

$$m\dot{\vec{v}} = \vec{F}_{ext} + \eta\left[\vec{v}(t-2a/c) - \vec{v}(t)\right] \tag{1}$$

here a - radius of the sphere, $\eta = \frac{Q^2}{3ca^2}$, $\vec{v} = d\vec{R}/dt$, $\vec{R}$ - coordinate of the center of the shell, $\vec{F}_{ext}$ - some external force. The term in RHS of eq.(1), proportional to η, describes the effect of radiation reaction (effect of retardation).

This model is a good tool to consider effects of radiation reaction, free of problems of classical point-like Lorentz-Dirac description (see, for ex.[3]).

A.

In this paper we consider Sommerfeld particle in the role of the Brown particle, i.e. particle (with radiation reaction), moving in a stochastic path under action of some external stochastic force $\vec{F}_{stoch}$.

For simplicity we take that:

(1) the viscosity of the surrounding medium is zero;

(2) the statistical average of $\vec{F}_{stoch}$ - $\vec{F}_0$ - is non zero and constant in time t;

(3) particle moves in one dimension.

Under these assumptions the Langevin equation for Sommerfeld particle takes the form

$$m\dot{v} = F_{stoch} + \eta\left[v(t-2a/c) - v(t)\right] \tag{2}$$

or for statistical average value $< v >$ of v

$$m< \dot{v} > = < F_{stoch} > + \eta\left[< v > (t-2a/c) - < v > (t)\right] \tag{3}$$

For dimensionless variables $y = < v > /M, \quad x = ct/M$ (M -scale factor) equation (3) takes the form

$$\dot{y} = f + \gamma \cdot \left[y(x-\delta) - y(x)\right] \tag{4}$$

here

$$\gamma = \frac{Q^2 M}{3a^2 mc^2}, \quad f = \frac{F_0 M}{mc^2}, \quad \delta = \frac{2a}{M}$$

Classical analog of equation (4) for Brown point particle one can get taking $\gamma = 0$ in (4).

Equation (4) for $f = const$ has the exact solution

$$y = y_0 + k \cdot x \tag{5}$$

with $k = f/(1+\gamma\delta)$ and initial velocity y_0.

Following the theory of Brownian motion (there are many textbooks on Brownian motion, see, for ex., [4]), the dispersion $D = < (v - < v >)^2 >$ for surrounding medium without viscosity can be function of time: $D = D(t)$. The form of $D(t)$ strongly depends on the form of correlation function of stochastic force F_{stoch} and its concrete realization in computer program (if correlation function is compact enough, then $D(t)$ for "not very large moments of time t" is proportional to time: $D(t) \sim t$ - Einstein formula for

Brownian motion). If the dependence $D = D(t)$ is known, one can try to find the form of correlation function, but this is not our goal.

Do not going into details, we can say that time average of D with respect to the whole time of "observation" T must be proportional (in dimensionless variables) to the effective temperature θ and inverse to the mass of particle:

$$\bar{D} \sim \frac{\theta}{mc^2}. \tag{6}$$

Solution (5) describes motion of charged particle with constant acceleration. This result seems to be unusual.

Indeed, following classical electrodynamics, particle with acceleration must radiate. Then one could expect that radiation, due to energy loss, leads to radiation damping of particle motion.

Instead, we see that after statistical averaging, radiation reaction in Sommerfeld form only changes the value of the effective force, acting on particle, making it smaller.

One can interpret this in the following way. Radiation reaction F_{rad} in Sommerfeld model is

$$F_{rad} = \gamma \cdot [y(x-\delta) - y(x)]$$

on trajectory (5) it is nonzero and equals to $-\gamma\delta k$. If δ - is small we can expand the force F_{rad} in powers of δ:

$$F_{rad} = -k\delta\dot{y} + k/2(\delta)^2\ddot{y} + ... \tag{7}$$

First term in (7) is the effective electromagnetic mass of the particle, multiplied by acceleration with sign minus. On trajectory (5) this term equals to $-\gamma\delta k$. Second term in (7) is the radiation force in classical Lorentz form - on trajectory (7) it is zero. Thus radiation effects after statistical averaging lead only to change in effective particle mass (mechanic + electromagnetic) - it becomes greater and the particle becomes more "inertial" (more "retarded").

It looks like the energy of self-electromagnetic field of a charged particle does not radiate away, the "electromagnetic fur-coat" does not get thin, thus particle becomes "heavier" (in comparison with particle of zero charge).

This effect can also make the dispersion (6), i.e. the effective Brownian temperature smaller.

B.

Dispersion (6), also as the particle motion, we investigated numerically. The particle mass m and size a we take close to that of classical electron, this yields $\gamma\delta = 1$ in (4).

The stochastic force F_{stoch} we extract step by step from the known procedure (see, for ex.,[5]):

$$\phi_{n+1} = \{K \cdot \phi_n\} \, ;$$

here brackets {...} denotes fractional part of ...,

and

$$(F_{stoch})_n \cdot (M/mc^2) = 10^{+3} \cdot (\phi_n - \phi_0)$$

with $K = 100000/3$ and $\phi_0 = 0.5007645$.

The results of numerical calculations can be summed in the following way:

(1) Sommerfeld particle in the role of the Brown particle has the effective temperature θ_S lower, than that for classical Brown particle θ_B: $\theta_S < \theta_B$.

(2) The more is the retardation (i.e. the greater is γ in (4)), the greater is the difference between θ_B and θ_S.

(3) The acceleration k of the average velocity of Sommerfeld particle is smaller than that of classical Brown particle without electric charge.

These results are illustrated in Fig. 1, where $y_0 = 0.1$; $\gamma = 2000.0$; $\delta = 2a/M = 1/2000.0$; observation "time" is 10.0005, with respect to it the average value of stochastic force is $f = 4.881996 \cdot 10^{-4}$ and the average dispersions are $\bar{D}_S = 4.4359 \cdot 10^{-8}$, $\bar{D}_B = 4.8496 \cdot 10^{-8}$, i.e. $\theta_S < \theta_B$.

The upper curve - is the path of classical Brown particle (horizontal axis - is "time" x), the lower - is the path of Sommerfeld particle in the role of Brown particle.

The acceleration of the average velocities for these particles differs by multiplier 2, as the consequence of the exact solution (5) (for Brown particle we have $\gamma = 0$, and $k = f$, for Sommerfeld - $\gamma\delta = 1$ and $k = f/2$).

These results confirm numerically our considerations, made before.

REFERENCES

1. A.Sommerfeld, Gottingen Nachrichten, 29 (1904), 363 (1904), 201 (1905).

2. L.Page, Phys.Rev., 11, 377 (1918). T.Erber, Fortschr. Phys., 9, 343 (1961). P.Pearle in "Electromagnetism",ed. D.Tepliz, (Plenum, N.Y., 1982), p.211. A.Yaghjian, "Relativistic Dynamics of a Charged Sphere". Lecture Notes in Physics, 11 (Springer-Verlag, Berlin, 1992). F.Rohrlich, Phys.Rev.D, 60, 084017 (1999).

3. Alexander A.Vlasov, physics/9905050, physics/9911059, physics/0004026.

4. I.A.Kvasnikov, "Thermodynamics and statistical physics. Part 2". Moscow, Moscow State University, 1987.

5. R.Z.Sagdeev, G.M.Zaslavsky, "Introduction to nonlinear physics". Moscow, Nauka, 1988.


1.05e-1
1.04e-1
1.03e-1
1.02e-1
1.01e-1
9.97e-2
0.00e0
1.00e0
2.00e0
3.00e0
4.00e0
5.00e0
6.00e0
7.00e0
8.00e0
9.00e0
1.00e1

Fig. 1